

## MATHEMATICIANS IN OUR LIVES

With the support of



11-12 years olds

## INTRODUCTION

As part of the *Mathematicians in Our Lives* programme, the Irish Mathematical Trust (IMT) has developed this package to celebrate the life and legacy of Sir William Rowan Hamilton. The lesson is structured in four sections: After a short review of Hamilton's life and achievements, we will focus on the main areas of his work, with sections on Geometrical Optics, Graph Theory, and Quaternions. Each section discusses the fundamental questions in the area and some of Hamilton's contributions, and includes a set of suggestions for class discussions, games and hands-on exercises. The lesson plan is designed so that you may extract sections to teach or to use the content to build lessons around the information provided. We hope that you enjoy this exploration of the brilliant mind that was Sir William Rowan Hamilton's.

Objectives:

- To introduce William Rowan Hamilton as a person and as a Mathematician.
- To explain the basics of the Laws of Optics, Graph Theory, Quaternions.
- To illustrate the rich interplay between Algebra and Geometry through examples from Optics, Graphs, Complex numbers and Quaternions.
- To solve games, practical tasks and logical exercises on the topics above.

Required:

- One copy of [Hamilton Museum Circuits](#) for each student.
- One [Quaternion Ball kit](#), scissors and stapler or sellotape for group of 2-3 students.
- One copy of the worksheet per student.

Lesson time: 1-3 lessons of 40 min each.

## WHO IS WILLIAM ROWAN HAMILTON?


Sir William Rowan Hamilton (1805-1865), one of Ireland's most famous scientists, was a distinguished mathematician, physicist and astronomer. He made a large number of important contributions to Optics, Mechanics, Graph Theory and Algebra. Many notable concepts in physics take their name from him; like the Hamiltonian function and Hamiltonian mechanics, while in Algebra, his best-known discovery is of the Quaternion number system.

## HIS LIFE

- *The story of a childhood at the same time normal and exceptional. Invite your students for their opinions on whether there was a connection between the early training in languages and the later proficiency in mathematics?*
- *Starting from Hamilton's story, invite students to discuss how initial challenge and defeat can influence a person's future career.*
- *Ask your students if they ever visited an observatory. Give a short description of one. What connections can be found between a job at an observatory and research in mathematics?*
- *Propose further historical investigation: Compare the lives of George Boole and William Rowan Hamilton. Did they live in Ireland at the same time? Did their lives/work intersect?*

William Rowan Hamilton was born in Dublin on 4<sup>th</sup> August, 1805. Judging by all his academic exploits at an early age, you wouldn't believe that Hamilton was a healthy boy who loved swimming, nature and jolly gatherings of friends. By the age of 13, nature walks brought out his enthusiasm in the form of poetry in at least 13 languages (Latin, Greek, Persian, Hebrew, Arabic, Sanskrit and others). His education was in the hands of his uncle, an accomplished linguist; Hamilton's mother and father had both died by the time he was 14.

The young Hamilton's first recorded mathematical adventure was a contest that pitted him against another child prodigy, the American "calculating boy" Zerah Colburn (unfortunately, Hamilton lost). Once Hamilton's curiosity about mathematics was ignited, its fire spread rapidly in his imagination. He entered Trinity College Dublin to study both classics and mathematics – achieving the highest honours in both - but he was more and more attracted by the later. He started blending algebra and geometry to study the laws that explain how light moves. He hadn't yet completed his studies when he presented his great work "*Theory of Systems of Rays*" to the Irish Academy (April, 1827). In the same year, before he had time to finish the final exams, he was appointed professor of astronomy, ahead of some well-established astronomers, and despite the fact that he hadn't even applied for the job!



Hamilton worked at the Dunsink Observatory till the end of his life. This was a rich and layered life, with many friends among poets as well as scientists.

## HIS WORK – AN OVERVIEW

- *Outline the three main areas of Hamilton's work which will be investigated in this lesson: Optics; Graph Theory; Quaternions.*
- *Discuss practical applications of Hamilton's work: conical refraction; the transition from the Hamiltonian mechanics to quantum mechanics and its uses in modern life; the use of quaternion in describing 3D rotations for airplane/space-ship flights and for computer games. See more resources at the end of this document.*

Hamilton started his scientific work out of curiosity about optics: the laws that explain how light travels through different media like air, water and glass, and how it reacts to obstacles or other changes. In his work *“Theory of Systems of Rays”* and its supplements, he devised the idea of characteristic function: a tool for measuring the time it would take light to travel along various paths, in terms of the start and end coordinates. This allowed him to explain the laws of optics based on the principle that *light always chooses the fastest path* (the *Minimum Principle*).

This brought him to spectacular and unexpected predictions about lights’ behaviour. For example, people before him had observed one ray splitting into two or three when passing through a crystal, but Hamilton discovered that in certain cases there would be an infinite number, a cone of refracted rays – which was confirmed by experiments and won him a Royal Medal in Physics.

Even though your laser has emitted just one line beam, from the other side of certain crystals you see it as a ring of light:





Fig. 1. Transformation of laser beam by conical refraction in a biaxial crystal.

Hamilton’s work in geometrical optics fit in well with the new treatment of mechanics developed by J. L. Lagrange (1736-1813), but Hamilton brought a simplicity and clarity which allowed him to carry over all of his methods effortlessly to the most general problems of mechanics.

Almost one hundred years after Hamilton presented his work to the Royal Irish Academy, his methods were found to be just what was needed for the creation of quantum mechanics in 1925-1926, which has in turn brought us the marvels of the digital world.

In his later life Hamilton became more and more intrigued by the interplay between algebra and geometry. This led him to the discovery of the quaternions, a four-dimensional extension of complex numbers determined by the equations  $i^2 = j^2 = k^2 = ijk = -1$  (which he famously carved into the side of Broom Bridge in Dublin). He spent the greater part of the rest of his life studying the quaternions and their properties, putting forward applications in the study of rotations that are used in aero- and astronautics to this day.




In graph theory he introduced the notions of Hamiltonian paths and circuits while searching for a closed path along the edges of a dodecahedron that visits each vertex exactly once. These ideas generate theorems to this day.

## OPTICS


- *A short introduction into Optics through the example of light refraction.*
- *Encourage your students to get involved in a class discussion on the Minimum Principle:*
  - ◆ *How does it apply to Refraction in Optics?*
  - ◆ *Why do they think the principle holds true?*
  - ◆ *How did Hamilton apply it and how does it relate with the Google Maps directions app?*
- *You may organize the students in teams of 2-3 and let them work on choosing the best paths between two points when the medium of propagation changes.*

For millennia, people have been attracted to the night sky and the movement of stars – the main questions of astronomy. This is how Galileo Galilei (1564-1642) had come to invent the telescope in 1609, by cunningly exploiting a property of light called refraction.



Since then, the best minds of their time tried to find the true explanation for the refraction of light.

For example, when passing from water to air, the light ray bends. This is why our minds get tricked into perceiving a fish as closer to the surface than it is. Indeed, on the way from the fish to our eye, the light ray had bent, but our minds still think it's straight. So in our mind we "see" the fish in an imaginary position along a straight line, instead of its real position lower down.



## REFRACTION AND THE MINIMUM PRINCIPLE

**Class Discussion:** So, what causes the light to bend when passing from water to air?

**Answer:** The light travels faster through air than through water.


**Class Discussion:** Why do you think the light travels faster through air than through water?

**Answer:** If we could look at air and water through a powerful microscope, we'd see that they are made of little pieces we call molecules. The air is made of molecules of different gases, which keep apart from each other. The water molecules are more crowded. When the light hits a molecule, this shoves the light out of its path a little. Imagine getting shoved every other step – this is bound to slow you down.

**Class Exercise:** Plan Your Trip! Normally, the fastest path between two points is a straight line - but not when you hit obstacles, which cause delays.

(a) In the picture here, count the number of dots that touch the path AB to find how much the traveller is slowed down. The top dots represent gas molecules in the air. The lower dots are water molecules.

(b) Now try to plan a better trip from A to B: Choose a point C on the black separating line, connect it to both A and B by straight line segments, and count the total number of dots you crossed. Is it more or less than on the path AB?




Report your result and explain your strategy. The student who crossed the fewest dots has found the *fastest path* and wins.

**Sample Answer:** The red path is an example of a faster trip:

AOB hits 13 dots.

The red path hits about 8 dots. It is chosen so as to make the trip through water shorter, without making the trip through air too long.



Inspired by the ancient work of Heron of Alexandria (c. 10-70 A.D.), the French mathematician Pierre Fermat (1601-1665) came up with a *Minimum Principle*, which basically states:

*The light ray always travels along the fastest path.*



**Class Discussion:** How does a light ray plan its journey?

If you don't suppose that Light is as thoughtful as a human being, you might think there is something funny with the idea that the Light knows where it wants to go and plans the whole trajectory in advance.

What's going on? How does the *Minimum Principle* make sense?

**Answer:** First off, distances; when Hamilton and others applied the Minimum Principle, they were thinking of light travelling extremely small distances, so that you might argue the planning wouldn't require that much foresight. Secondly, we might think of the *Minimum Principle* more as an example of *observer effect*: we think that that's how light behaves because this behaviour is the only one that we can observe. Consider this: a light source usually sends rays in many directions. However, the light rays that start at A in a "wrong" direction will meet more and more obstacles that jolt and deflect them further away from the target. Thus their chances of hitting the target B (where we patiently await) become so small that we can hardly notice any such rays reaching the destination. In fact, our eye is not trained to notice such small effects.

On the other hand, finer measuring instruments may detect "stray" rays, with various frequencies. This kind of probabilistic thinking applied at extremely small scales inspired Quantum Mechanics, an area of Physics to which we owe much of our understanding of nature at atomic level, as well as semiconductor-based electronics, (computers, smartphones), optical cable telecommunication (the Internet), and other features of modern life.

**Exercise:** How to find the perfect path – with numbers.


In the Light SuperWorld, light rays travel on any paths they like. Three rays called Mr Simplex, Mrs Wiseman and James Bold, decide to go from a point A, found 100 meters above water, to a point B, found 100 meters below water. They are warned that travelling through water is slower, namely

- They can travel at a speed of 300 meters/second through air;
- But only 225 meters/second through water.

Mr Simplex decides to take a straight line from A to B, a total distance of 255 meters.

James Bold decides to go as much as possible through air, so he travels 187 m to point P, found exactly above B on the surface of the water, and then from P straight down to B.

Mrs Wiseman Ray makes some calculation and decides to go about 141.5 meters through the air, heading straight for a point W on the water surface, and then travels about 115.5 meters through the water, from W to B.



Which Ray gets to the destination fastest? Can you intuitively explain why?

**Answer:** Using  $time = distance/speed$  separately through air and water, we get:

For Mr Simples:  $t_{air} + t_{water} = \frac{127.5}{300} + \frac{127.5}{225} = 0.992 \text{ s.}$

For James Bold:  $t_{air} + t_{water} = 187/300 + 100/225 = 0.623 + 0.444 = 1.068 \text{ s.}$

For Mrs Wiseman:  $t_{air} + t_{water} = 141.5/300 + 115.5/225 = 0.47167 + 0.51333 = 0.985 \text{ s.}$

As light is slower through water, it makes sense to try to shorten the distance travelled through water, even if this makes the trip through air a little longer, too. This is the strategy that both Mrs Wiseman and James Bold took. However, James Bold lengthened the total distance too much. Whereas Mrs Wiseman tried to find a balance between the time spent in the air and the time spent in the water.

*Play hands-on* with refraction angles [here](#):


[www.physicsclassroom.com/Physics-Interactives/Refraction-and-Lenses/Refraction/Refraction-Interactive](http://www.physicsclassroom.com/Physics-Interactives/Refraction-and-Lenses/Refraction/Refraction-Interactive)

### SO... WHAT ABOUT HAMILTON?

Hamilton’s great insight was a smart way to calculate how fast paths are. He designed his calculation as a function of the coordinates of both the starting point and the target. In many more complicated problems, this viewpoint brought clarity and simplifications.

Today, Google Maps works much like Hamilton’s characteristic functions:

- It can take as inputs your starting points and the desired destination
- It calculates the duration of each possible path
- And it highlights the fastest path for your convenience:



Google Maps’ is a simpler set-up than in Optics, because cities have a finite numbers of possible paths. A map can be modelled mathematically by a graph whose points are all addresses and edges are the streets between them. You might not be surprised to find that Hamilton was also interested in graphs and their properties, and he has a special type of graphs named in his honour!

But wait! Guess what? All the data used by Google Maps in its algorithms was gathered by the Global Positioning System, a network of satellites around the Earth. To determine distances, they


use atomic clocks designed using principles of Quantum Mechanics – which profited much from Hamilton’s mathematical formulations.

## GRAPH THEORY

- *Introduce the area of Graph Theory using a famous puzzle.*
- *Define the notions of Hamiltonian paths, cycles and graphs.*
- *Play some games based on identifying Hamiltonian cycles, and work through some applications.*


### THE SEVEN BRIDGES OF KÖNIGSBERG

The oldest and most famous use of graphs to describe travel around cities comes from the city of Königsberg (now the Russian city Kaliningrad). During the time of the Swiss mathematician Leonhard Euler (1707 – 1783), this was a Prussian city that lay on the Pregel River. A small island was located in the middle of the river at the city centre, and the 4 separate land masses were joined by seven bridges as shown.



The story goes that the people of the city invented a game, whereby they had to try to find a route through the city centre that crossed each of the seven bridges exactly once (without necessarily starting and finishing at the same point). Of course, going half-way across a bridge and turning back was not allowed, and neither was swimming, jumping the gap or running down the bank to look for an eighth bridge or hovercraft. Provided these rules were obeyed, it seemed that no-one could find a solution. Can you? Give it a go!

...but don't spend too long at it, because it's actually impossible. In fact, Leonhard Euler proved mathematically that no solution exists, and in doing so kick-started graph theory. Euler discarded most of the beautiful features of the 4 land areas in the city, and represented each area by one node (dot). He could then focus on the bridges and represented them as edges (curved lines connecting the dots):




He didn't care about correct sizes and locations, all he was interested in is what connects to what. Can you see the connection between the two pictures above? Next, Euler noticed that, except for the start and end points of your trip, whenever you enter a node by an edge, you also leave it by an edge, i.e. edges are used in pairs, and as a result, there must be an even number of edges connected to each vertex that isn't the start or end point. But you'll notice that all of the nodes here have an odd number of edges. Because of this, Euler concluded that you can't win the game.




As well as being a fun puzzle, this led to the mathematical field of graph theory. For the first time, a real-world situation had been replaced by an equivalent *graph*, the problem had been solved in this abstract setting, and the result translated back into the real world. A *graph* in graph theory is just a number of nodes connected by edges. We're now going to see how Hamilton contributed to the study of graphs.

### HAMILTONIAN PATHS

What we were looking for in the last section was an *Eulerian Path*, or a path through a graph that visits each edge exactly once. Hamilton was fascinated by shapes like the dodecahedron (aka football) and he started searching for a way along the edges that visits each vertex (node) exactly once. He didn't care about using all edges.



A path that visits each node of a graph once is now called a *Hamiltonian Path*, while a *Hamiltonian Cycle* is a Hamiltonian path that starts and finishes at the same point. The task of finding a Hamiltonian cycle on the edge-graph of a regular dodecahedron is called Hamilton's game or the icosian game. Let's give it a go!




First of all, let's change it to a 2-dimensional problem. Like in the last section, we don't care about the distances between nodes, provided the same things are connected to each other. So, we can "flatten out" the dodecahedron into a 2-D graph:

*Exercise:* Find a Hamiltonian cycle in this graph:

*Sample solution:*

Hamilton invented a new mathematical method called icosian calculus and tried to make this into a commercial product. However this ended in failure because the number of solutions people could find was small enough and they became bored of it too quickly.



The notion of Hamiltonian paths and circuits is the most interesting aspect of this story, and is an important part of graph theory to this day.

By Christoph Sommer (Own work) [[GFDL](#), [CC-BY-SA-3.0](#) or [CC BY-SA 2.5-2.0-1.0](#)], via Wikimedia Commons

### HAMILTONIAN GRAPHS


A graph is called Hamiltonian if it has at least one Hamiltonian Cycle in it.




### Application: Knight's Tour and the T.S. Problem

As you can imagine, there are so many different kinds of Hamiltonian path problems. A well-known example is the Knight's Tour in a chessboard:

Making a knight's tour involves moving a knight around a chessboard, landing at each square exactly once (no more nor less often). Knights can only move in "L-shapes", i.e. one square horizontally and two vertically or two squares horizontally and one vertically.



An "open" tour is one that doesn't start where it began; a "closed" tour does. It mightn't seem like this has anything to do with Hamiltonian paths, but think about it: what if we joined each square (using lines) to all squares the knight could move to in one turn? We'd get a graph, with the squares acting as the nodes and the lines as the edges. In fact, it would look like this:




Then, the Knight's Tour game becomes a matter of finding a Hamiltonian path or cycle in this graph. As you can imagine, there are loads of different tours that can be constructed. It is still a very difficult problem, however. It can't really be solved by brute force (listing all possible paths in

the graph and picking out the ones that are Hamiltonian) because there are around  $4 \times 10^{51}$  Hamiltonian paths that a knight can take (that's a fairly big number. In fact, if it took you a minute to check each path, it would take about 7.5 billion trillion trillion trillion years in total). There are, however, some algorithms that can produce results. A useful one is Warnsdorff's rule (a "heuristic" algorithm), which tells you that, when choosing your next move, you should always take the option with the lowest number of possible moves. If you ever find that two or more options share this lowest number, you should just pick one of them randomly (there are methods for determining which one to choose, but they're quite complicated).

[Here's](#) a cool Knight's Tour game which can be played online. It's easier to use than the traditional chessboard because it keeps track of the moves you've already made and only permits legal knight moves.

An extension of the Hamiltonian circuit problem is the Travelling Salesman Problem.

**Puzzle:** Imagine you are a salesperson who travels around a country selling your product in big cities. Some of the cities are linked by highways, while others aren't, and every highway-link between two cities has an associated distance. You want to visit every city exactly once and finish where you started, while at the same time ensuring you travel the smallest distance possible. Which path do you take?



Basically, the cities and highways can just be viewed as a graph in which the distance between nodes actually counts. The problem is simply to find the Hamiltonian circuit with the shortest length:

**Solution:**

When we calculate the length of the circuit, it doesn't matter where we start from. So we'll always start at A.

| Hamilton Circuit | Length         |
|------------------|----------------|
| ABCEDA           | $2+4+3+3+8=20$ |
| ABCDEA           | $2+4+7+3+6=22$ |
| ABDCEA           | $2+3+7+3+6=21$ |
| ADBCEA           | $3+3+4+3+6=19$ |

And the winner is ADBCEA.

This actually has countless applications in the real world. There are the obvious ones like deciding how a postman should most efficiently deliver mail, but other uses include determining how best to time online advertisements, place vanes on an aircraft turbine and wire a computer. It is in fact one of the most intensely-studied problems in optimisation.

## ALGEBRA AND GEOMETRY: FROM REAL NUMBERS TO COMPLEX NUMBERS AND QUATERNIONS:

- *Look at the Algebra with real numbers as a way to describe movements along a line.*
- *Introduce Complex Numbers as points in the plane, and operations with complex numbers as movements on the plane.*
- *Introduce Quaternion Algebra with the hands-on Quaternion Ball tool.*
- *Perform rotations in 3D using Quaternions*


Mathematics is a story spanning thousands of years, with hundreds of characters, both human and mathematical. It is a story too long for anyone to hear the whole of, but many people spend their lives listening. Today we will give a little bit of this story and our main characters will be an Irish mathematician, William Rowan Hamilton, and a new number system called the quaternions. But what are the quaternions and more importantly, why should we care about them? Hamilton came about the idea of quaternions as a way to represent rotations in a three-dimensional space.

## NUMBERS ON THE REAL LINE


In order to understand these fully, we need to remember how numbers and operations can be thought of in practical (and geometric) terms:

Real numbers = Points on a line  
Operations with numbers = movements along the line

For example,  $+3$  means we skip three unit steps to the *right*, starting from wherever we are.




Here we use the symbol  $x$  to describe a number that we might not know from the beginning, and whose value we might find out later. If our walk along the number line led us to 5, we write this as  $x + 3 = 5$ . To find  $x$ , we move back 3 spaces:  $x = 5 - 3 = 2$ .



Thus  $-3$  means moving 3 steps to the *left*. This is the type of thinking that the Persian mathematician *al-Khwārizmī* (780 – 850) described by the Arabic word *al-jabr* (reunion of broken parts), which is the origin of the well-known word *Algebra*.

Algebra uses symbols to describe numbers. This allows us to make general statements like this one (where  $a$  stands for any positive number):

$+a$  is a move of length  $a$  to the right, while  $-a$  is a move of length  $a$  to the left.



From the picture we see that  $-a$  is the result of the *reflection* of  $a$  across 0. Since

$-a = (-1) \times a$  we can thus give a geometric meaning to the multiplication by  $-1$ :

Geometrically, multiplication by  $(-1)$  is the *reflection* across 0.


$(-1) \times$  has the effect that every number to the left of 0 gets moved to the right and every number to the right of 0 gets moved to the left.

**Exercise:** Use reflection to explain why multiplying two negative numbers gives you a positive number.

**Solution:** As  $-b = b \times (-1)$  and  $-c = c \times (-1)$ , it is enough that multiplication by  $(-1) \times (-1)$  means reflecting twice, which takes any number back where it was originally. So  $(-1) \times (-1) = 1$ .

**Exercise:** Work out how to reflect around a number other than 0. Let's say that we have a number  $A$  on the line and another number  $x$ . Write an equation for the reflection of  $x$  through  $A$ . It should be an expression in  $x$  and  $A$ .

**Solution:**



Geometrically, we can move the whole line to the left by  $A$ . Then  $x$  becomes  $x - A$  and  $A$  becomes 0. Then reflecting through 0 sends  $x - A$  to  $-(x - A) = A - x$ . Now we need to move the line back to the right by  $A$ . This makes  $A - x$  into  $2A - x$ .


## COMPLEX NUMBERS

As often happens with stories, we must now skip ahead in time. We come to the second major characters in our story, the complex numbers. Armed with all the numbers, operations and symbols they could put on a line, mathematicians could now solve so many different equations that they even starting looking into impossible ones, like this one:

$$x^2 + 1 = 0$$

which is the same as  $x^2 = -1$  or equivalently  $x = \sqrt{-1}$  or  $-\sqrt{-1}$ . But taking the square root of a negative number seemed totally impossible for a long time. The first person who dared mention it was the sixteenth century Italian mathematician Cardano, who called such a number  $x$  meaningless, fictitious, and imaginary. From here on, this number was denoted by  $i$  from “imaginary”.

“For well over two centuries after imaginary numbers broke into the domain of mathematics they remained enveloped by a veil of mystery and incredibility until finally they were given a simple geometrical interpretation by two amateur mathematicians: a Norwegian surveyor by the name of Wessel and a Parisian bookkeeper, Robert Argand”. According to their interpretation a complex number, as for example  $3 + 4i$ , may be represented as in the Figure here in which 3 corresponds to the horizontal distance, and 4 to the vertical. (George Gamow, “One, two, three ... infinity”).



Looking at this geometrically, we now see


Complex numbers = Points on the plane

and hence we would expect that


Operations with complex numbers = movements on the plane.

**Products:**

When we multiply a real number, say 4, representing a point on the horizontal axis, by the imaginary unit  $i$  we obtain the purely imaginary number  $4i$ , which must be plotted on the vertical axis.



Multiplication by  $i$  is the same as a counter-clockwise rotation by a right angle around 0.



We can see that rotating by  $90^\circ$  twice corresponds to a rotation by  $180^\circ$ , which is the same as the reflection through the origin O:

$$i \cdot i \cdot 4 = -4$$

or equivalently  $i^2 = -1$ . The equation that had baffled mathematicians for centuries now has a very nice geometric interpretation.

(Recall that  $i \cdot$  is the rotation by  $90^\circ$  while  $(-1) \cdot$  is the reflection through 0.)

By the same means


Multiplication by  $-i$  is the same as a clockwise rotation by a right angle around 0.

**Reflections:**

Another way to move things in the plane is to reflect across lines. Reflecting a point across the line means drawing a segment through  $P$  which is perpendicularly bisected by the line.

For the reflection of  $z = a + ib$  across the horizontal and the vertical axis, respectively we have:


Reflection of  $z = a + bi$  across the real axis gives  $a - ib$ . This is called the conjugate of  $z$  and is denoted by  $\bar{z}$ . Reflection across the imaginary axis gives  $-a + ib$ .



*Exercise: Successive Reflections.*

Reflections are important in many ways. For example, take two lines  $l$  and  $k$  meeting at  $O$  and with an angle  $\alpha = 30^\circ$  between them. Reflect a point  $P$  through  $l$  and then  $k$  successively to get  $P'$  and then  $P''$ .

If  $\angle MON = 30^\circ$  and  $\angle POM = 10^\circ$ , find  $\angle POP''$ .



**Solution:** reflection means that  $\triangle POM \cong \triangle P'OM$  and  $\triangle P'ON \cong \triangle P''ON$  hence  $|PO| = |P'O| = |P''O|$  and  $\angle POM = \angle P'OM$  and  $\angle P'ON = \angle P''ON$ . Hence by summing up :  
 $\angle POP'' = \angle POP' + \angle P'OP'' = 2\angle P'OM + 2\angle P''ON = 2\angle MON = 2\alpha$ .

Two successive reflections across two lines through  $O$  amounts to a rotation by double the angle between the two lines  
 (clockwise or counter-clockwise depending on which line you reflected across first)

**THE QUATERNIONS**


- Introduce the Quaternions as a four-dimensional system of imaginary numbers.
- Describe addition and multiplication of Quaternions, with the use of a geometric visualisation.
- Consider some applications of the Quaternions.

Just like complex numbers numbers  $a + bi$  represent points in the plane and are made of a pair of real numbers  $(a, b)$ , we can represent a point in the 3-dimensional space by a triplet  $(x, y, z)$ . Hamilton was fascinated by the discovery that multiplication represents rotation in the complex plane, and he wanted to do the same in 3D. The problem of finding an algebra of triples  $(\alpha, \beta, \gamma)$  to describe the geometry of vectors in three dimensional (3D) space haunted him for at least fifteen years.

*“Every morning in the early part of the above-cited month [October 1843], on my coming down to breakfast, your (then) little brother William Edwin, and yourself, used to ask me, “Well, Papa, can you multiply triplets”? Whereto I was always obliged to reply, with a sad shake of the head: “No, I can only add and subtract them.” W R Hamilton in a letter dated August 5, 1865 to his son A H Hamilton [1].*

In 1843, Hamilton found an ingenious way around his problem. The solution famously came to him as he was walking along the Royal Canal in Dublin with his wife on 16<sup>th</sup> October (now called Hamilton day)- he suddenly realised that the answer lay in numbers with four components instead of three. In his excitement, he promptly used his penknife to carve the solution equations into the side of nearby Broom bridge:

$$i^2 = j^2 = k^2 = ijk = -1$$




Plaque on Broom bridge. Wikimedia Commons.




This was the birth of the quaternions, which we represent by  $\mathbb{H}$  in honour of Hamilton. Let's look more closely at what we've just written.

First, Hamilton decided that he could have not just 1, but 3 "imaginary" axes, each with its own unit:  $i$ ,  $j$  and  $k$  are what we call the quaternion units. They form the building blocks of the quaternions. Notice that they all satisfy  $i^2 = j^2 = k^2 = -1$ , so in fact we now have 4 axes: three for  $i$ ,  $j$  and  $k$ , and a fourth one for the real numbers.

To help us better grasp the consequences of the rules above, we need to play with the symbols  $i$ ,  $j$  and  $k$  and in particular, to understand their products. Luckily, we have a handy toy available to help us with this task. It is based on the following ball in the 3D quaternion space. As you can see, the axes are marked by the units  $i$ ,  $j$ ,  $k$  on one side, and  $-i$ ,  $-j$ ,  $-k$  on the other side. On each of the three coordinate planes marked by circles, multiplication  $\cdot i$ ,  $\cdot j$ ,  $\cdot k$  represents a rotation. Indeed, each circle lies on a plane very similar to the complex plane we're met earlier.



If you haven't already done so, please print out and assemble the Quaternion Ball learning tool by clicking [here](#). We will use it to play with quaternion right multiplication. As you can see, the Ball is made of three discs that intersect at right angles, with red circles on each disc. We find that tracing a finger along these circles while carrying out this exercise is helpful. Tracing out a quarter circle in the same direction as the arrow on it corresponds to right the quaternion unit ( $i$ ,  $j$  or  $k$ ) printed next to the circle. Tracing opposite to the arrow's direction corresponds to the negative of the unit. Tracing out a number of quarter arcs in sequence corresponds to each of the units traced out written down next to each other in the same order: to the right, for example, I trace out  $i$  and then  $-j$ , which matches the multiplication  $i(-j) = -ij$ . Any other path that takes you from the same starting point to the same finish gives an equal answer: here, I could also have taken  $-k$  to get to the same point, so I know now that  $-ij = -k$ , or  $ij = k$ . Let everyone in the class please copy the table just below onto a sheet of paper, and using the



Quaternion Ball as we've described, fill out all the missing entries. Note: Multiplication by 1 does not figure on the Quaternion Ball because it represents staying in place: no change.

| $\times$        | <b>1</b> | <b><i>i</i></b> | <b><i>j</i></b> | <b><i>k</i></b> |
|-----------------|----------|-----------------|-----------------|-----------------|
| <b>1</b>        |          |                 |                 |                 |
| <b><i>i</i></b> |          |                 | <b><i>k</i></b> |                 |
| <b><i>j</i></b> |          |                 |                 |                 |
| <b><i>k</i></b> |          |                 |                 |                 |

*Exercise:*

Fill in the table with the correct Hamilton products. For each box, its row represents the first number in the product, while its column is the second number.

For example, I've place ***k*** in row ***i*** and column ***j*** because  $ij = k$ .

*Solution:*

Once you've filled in the box, you will notice something kinda weird about it. Everybody knows that

| $\times$        | <b>1</b>        | <b><i>i</i></b>   | <b><i>j</i></b>   | <b><i>k</i></b>   |
|-----------------|-----------------|-------------------|-------------------|-------------------|
| <b>1</b>        | 1               | <b><i>i</i></b>   | <b><i>j</i></b>   | <b><i>k</i></b>   |
| <b><i>i</i></b> | <b><i>i</i></b> | -1                | <b><i>k</i></b>   | - <b><i>j</i></b> |
| <b><i>j</i></b> | <b><i>j</i></b> | - <b><i>k</i></b> | -1                | <b><i>i</i></b>   |
| <b><i>k</i></b> | <b><i>k</i></b> | <b><i>j</i></b>   | - <b><i>i</i></b> | -1                |

reversing the order of addition doesn't matter, don't they?


$$3 + 4 = 7 = 4 + 3, \text{ right?}$$

Similarly for multiplication:  $2 \times 5 = 10 = 5 \times 2$ .

However, we've just said that  $jk = i$  and  $kj = -i$ , which aren't equal. So, order **does** matter when it comes to the quaternions: that's just how they are. That's something to watch out for when dealing with them.

Note, however, that when you multiply just an imaginary and a real, order doesn't matter. For example,  $(8)(j) = (j)(8)$ .

### USING THE QUATERNIONS – ROTATIONS IN 3D




Aside from being a fascinating area of study in pure maths, the quaternions have at least one major application in the real world—mathematically representing rotations in three dimensions. There are other ways (like using rotation matrices) to do this, but the algebra is much simpler when using quaternions and certain problems that can arise with matrices are avoided. As a result, rotations are described by quaternions in multiple areas, including video game and movie animation, aircraft and spacecraft attitude control and robotics.


### Quaternions in Spaaace!

As we explained before, quaternions are used for controlling spacecraft attitude, so we're now going to look at an example of how this works. By the way, attitude is just a terms used in astronautics and


other areas that means orientation in 3-D. Now, let's think about a space shuttle in orbit above earth:



*Example:* Imagine the astronauts are having a tanning competition on board and want to turn the shuttle by  $90^\circ$  to its left so that it faces the sun, like this:



How does the shuttle carry out this command? To be mathematical, we say that we want to rotate the shuttle by  $90^\circ$  around a vertical axis:




*Answer:* Recall that multiplication by  $i$  represents a counter-clockwise rotation by  $90^\circ$  on the horizontal plane. This rotation is simple though; other rotations do not have formulas as simple.

Remember that Hamilton discovered quaternions on the way to explaining how to use multiplication to rotate objects in 3D. We will work in the 3D space whose 3 axes have units  $i, j$  and  $k$ .

Recall the Quaternion Ball we have played with earlier. First off, we should notice that although each on their own circle, the multiplications  $\cdot i$ ,  $\cdot j$  and  $\cdot k$  represent rotations by  $90^\circ$ , none of them represents a rotation on the entire 3D space.

Indeed,  $i^2 = j^2 = k^2 = -1$ , which is a plain number and does not even exist the 3D-space of  $i, j, k$  (whose points are all of the form  $xi + yj + zk$ ).

Thus trying to “rotate”  $i$  by  $i$  would land us in a 4<sup>th</sup> dimension altogether! Depending on your viewpoint, this is either very exciting or disappointing:



Product of quaternions is not rotation in 3D space.

But let's not despair. The product of quaternions has all kinds of nice geometric connections which we will explore.

*Exercise: Geometric Exploration:* Here's a nice transformation in 3D:


a) Calculate the products  $iji$ ,  $iki$  and  $iii$ . Now take a number and surround it by  $i$  and  $i$ , like this:  $i(ai + bj + ck)i$ . Plot your results in the 3D space. What do you notice? Can you describe this operation as a geometric transformation (movement) of the point  $p = ai + bj + ck$ ?

b) Try the same problem replacing  $i$  by  $j$ : Calculate  $jij$ ,  $jjj$ ,  $jkj$ . What do you notice?


*Solution:*  $iii = -i, iji = ki = j, iki = ij = k$ ,

which is the reflection across the plane of axes  $j$  and  $k$ . This is the plane perpendicular to  $i$ .


Similarly, surrounding a point  $p$  by  $j$ -s is the same as reflecting it across the plane perpendicular to the  $j$  axis.



1. Rotations in Euler angles can be defined like gimbal system with three circles



2. When all three circles are lined up, the whole system can only move in two dimensions from this configuration, this is a gimbal lock



3. Usage of quaternions can help to avoid such situations

<https://www.quora.com/What-are-some-real-life-applications-of-complex-numbers-in-engineering-and-practical-life>

Hamilton was delighted with his discovery of Quaternions, and anticipated many future uses. Indeed, Quaternions have found uses varying from the positioning of planes and of space shuttles to computer graphics. Hamilton's great contribution to Mathematics does not consist so much in the Quaternions themselves, as much in setting an example of reshaping laws and crossing boundaries of thought, from solving 3D problems by introducing 4 dimensions to the new non-commutative Algebra he designed. Today, non-commutative algebraic and geometric methods continue to play a large law in Mathematics and Theoretical Physics.

---

This lesson plan was prepared by Thomas Sheerin, Anca Mustata and Jacob Bennett-Woolf for the Irish Mathematical Trust programme Mathematicians in Our Lives. We have used a large number of resources. We most warmly recommend in particular:

- "One Two Three . . . Infinity: Facts and Speculations of Science" - book by: [George Gamow](#)
- Wolfram MathWorld <http://mathworld.wolfram.com/>
- "Men of Mathematics. The Lives and Achievements of the Great Mathematicians from Zeno to Poincare" – book by E.T. Bell
- <https://www.forbes.com/sites/chadorzel/2015/08/13/what-has-quantum-mechanics-ever-done-for-us/#1c65408e4046>
- <https://www.slideshare.net/edzontatualia/refraction-of-light-45755877>